Efforts of the American Cancer Society, other organizations and individuals to obtain attention both at the local and national levels were noted in many news stories, columns and editorials.

COURIER-JOURNAL
Louisville, Kentucky
September 3, 1962

Smoke Study Adequate, Doctor Says

Chicago, Sept. 2 (UPI)—A New York doctor says further amassing of scientific data linking cigarette smoking to lung cancer is not necessary.

"Studies already completed are adequate and are available for reevaluation if that is needed, said Dr. Howard C. Taylor in the September Journal of the American Medical Association.

"The evidence is far more precise, in fact, than that which has been accepted by physicians as adequate to take action against the causes of other plagues," Taylor wrote.

Evidence Cited

"There is the known presence in cigarette smoke of carcinogenic (cancer-causing) substances, the production of skin cancer in laboratory animals by smoke condensates, the predilection of pre-cancerous lesions for the lungs of heavy smokers, and the numerous epidemiological and statistical studies showing a constant parallelism between the number of cigarettes habitually smoked and the chance of developing lung cancer.

"Nevertheless," the doctor added, "the individual smoker does not relate the hazard to himself, either because the extent of the risk is not comprehended or the day of reckoning seems too remote."

THE CHARLOTTE OBSERVER
Charlotte, North Carolina
August 31, 1962

SAYS N.Y. DOCTOR

Cigarette-Cancer Tie 'Real'

CHICAGO — A New York doctor said Thursday that further amassing of scientific data linking cigarette smoking to lung cancer is not necessary.

"Studies already completed are adequate and are available for re-evaluation if that is needed," said Dr. Howard C. Taylor in the September Journal of the American Medical Association.

"The evidence is far more precise, in fact, than that which has been accepted by physicians as adequate to take action against the causes of other plagues," he wrote.

"There is the known presence in cigarette smoke of carcinogenic (cancer-causing) substances, the production of skin cancer in laboratory animals by smoke condensates, the predilection of pre-cancerous lesions for the lungs of heavy smokers and the numerous epidemiological and statistical studies showing a constant parallelism between the number of cigarettes habitually smoked and the chance of developing lung cancer.

"Nevertheless," the doctor said, "the individual smoker does not relate the hazard to himself, either because the extent of the risk is not comprehended or the day of reckoning seems too remote."

(Dr. Taylor is the chairman of the ACS Committee on Tobacco and Cancer, though neither J.A.M.A. nor news reports noted this.)

1005150620

1005150620 A

Source: https://www.industrydocuments.ucsf.edu/docs/nhyk0000